



July Forecast Update for Northwest Pacific Typhoon Activity in 2002

Issued: 11th July 2002

by Drs Paul Rockett and Mark Saunders
Benfield Greig Hazard Research Centre, UCL (University College London), UK

Forecast Summary

TSR anticipates Northwest Pacific typhoon and intense typhoon activity being 15-30% above the 10-year climate norm in 2002.

The TSR (Tropical Storm Risk) consortium early July forecast update continues to anticipate an active typhoon and intense typhoon season for the Northwest Pacific. Storm numbers are expected to be 15-30% higher than the 10-year climate norm. The forecast spans the full Northwest Pacific season from 1st January to 31st December 2002 and is based on data available through the end of June 2002. Our main predictor is the forecast anomaly in August-September Niño 4 sea surface temperature (SST) which we anticipate will be $0.88 \pm 0.24^\circ\text{C}$ warmer than normal this summer. A final monthly forecast update will be issued in early August 2002.

NW Pacific Total Numbers in 2002

		Intense Typhoons	Typhoons	Tropical Storms
TSR Forecast (\pm FE)	2002	11.8 (\pm 2.2)	19.2 (\pm 3.7)	28.6 (\pm 4.4)
10yr Climate Norm (\pm SD)	1992-2001	9.0 (\pm 3.1)	16.9 (\pm 4.3)	27.4 (\pm 4.6)
30yr Climate Norm (\pm SD)	1972-2001	8.2 (\pm 3.3)	16.4 (\pm 3.6)	26.3 (\pm 4.0)
Forecast Skill at this Lead	1987-2001	39%	16%	8%

Key: Intense Typhoon	=	1 Minute Sustained Wind > 95Kts	=	Hurricane Category 3 to 5
Typhoon	=	1 Minute Sustained Wind > 63Kts	=	Hurricane Category 1 to 5
Tropical Storm	=	1 Minute Sustained Wind > 33Kts		
SD	=	Standard Deviation		
FE (Forecast Error)	=	Standard Deviation of Errors in Simulated Real Time Forecasts 1992-2001		
Forecast Skill	=	Percentage Improvement over Running 10-year Prior Climate Norm from Simulated Real Time Forecasts 1987-2001		
Northwest Pacific	=	Northern Hemisphere Region West of 180°W Including the South China Sea. Any Tropical Cyclone (Irrespective of Where it Forms) Which Reaches Tropical Storm Strength Within this Region Counts as an Event.		

Key Predictor for 2002

The key factor behind our forecast for an active Northwest Pacific typhoon season in 2002 is the anticipated enhancing effect of warmer than average Niño 4 (150°W-160°E, 5°S-5°N) SST. Our forecast anomaly (1972-2001 climatology) for August-September 2002 Niño 4 SST is $0.88 \pm 0.24^\circ\text{C}$ (up from $0.53 \pm 0.29^\circ\text{C}$ last month). Forecast skill for this predictor at this lead is 58% (assessed using replicated real-time forecasts over the last 15 years).

Further Information

Further information on the TSR forecast methodology, the TSR simulated real-time forecast skill 1987-2001 as a function of lead time, and on TSR in general, may be obtained from the 'Extended Range Forecast for Northwest Pacific Typhoon Activity in 2002' document issued on the 6th March 2002. Our forecast models for typhoon and tropical storm numbers have changed this month with these numbers now being obtained by regression from intense typhoon numbers. As a result our forecast numbers for typhoons and tropical storms have fallen from last month while our forecast for intense typhoons has risen as expected from the raised forecast for August-September 2002 Niño 4 SST. A final monthly forecast update for the 2002 Northwest Pacific Typhoon season will be issued on the 6th August 2002.

Appendix - Predictions from Previous Months

NW Pacific Total Numbers 2002				
		Tropical Storms	Typhoons	Intense Typhoons
Average Number (\pm SD) (1992-2001)		27.4 (\pm 4.6)	16.9 (\pm 4.3)	9.0 (\pm 3.1)
Average Number (\pm SD) (1972-2001)		26.3 (\pm 4.0)	16.4 (\pm 3.6)	8.2 (\pm 3.3)
TSR Forecast (\pm FE)	11 July 2002	28.6 (\pm 4.4)	19.2 (\pm 3.7)	11.8 (\pm 2.2)
	7 June 2002	30.8 (\pm 4.5)	21.1 (\pm 3.5)	10.5 (\pm 2.2)
	7 May 2002	30.5 (\pm 4.6)	20.9 (\pm 3.4)	10.3 (\pm 2.2)
	5 Apr 2002	29.6 (\pm 5.0)	19.8 (\pm 4.1)	9.8 (\pm 2.6)
	6 Mar 2002	28.6 (\pm 4.8)	18.7 (\pm 4.0)	9.3 (\pm 2.5)

